Relationship Between the Hosoya Polynomial and the Edge-Hosoya Polynomial of Trees

Niko Tratnik[a], Petra Žigert Pleteršek[a,b]

[a] *Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia*
[b] *Faculty of Chemistry and Chemical Engineering, University of Maribor, Slovenia*

arXiv:1609.04182v1 [math.CO] 14 Sep 2016

Abstract

We prove the relationship between the Hosoya polynomial and the edge-Hosoya polynomial of trees. The connection between the edge-hyper-Wiener index and the edge-Hosoya polynomial is established. With these results we also prove formulas for the computation of the edge-Wiener index and the edge-hyper-Wiener index of trees using the Wiener index and the hyper-Wiener index. Moreover, the closed formulas are derived for a family of chemical trees called regular dendrimers.

Keywords: Hosoya polynomial, edge-Hosoya polynomial, tree, dendrimer, edge-hyper-Wiener index
2010 MSC: 92E10, 05C31, 05C05, 05C12

1. Introduction

The first distance-based topological index was the Wiener index introduced in 1947 by H. Wiener [1]. Later, in 1988 H. Hosoya [2] introduced some counting polynomials in chemistry, among them the Wiener polynomial, which is strongly connected to the Wiener index. Nowadays, it is known as the *Hosoya polynomial*. Another distance-based topological index, the hyper-Wiener index, was introduced in 1993 by M. Randić [3]. All these concepts found many applications in different fields, such as chemistry, biology, networks.

The Hosoya polynomial, the Wiener index, and the hyper-Wiener index are based on the distances between pairs of vertices in a graph, and similar concepts have been introduced for distances between pairs of edges under the names the edge-Hosoya polynomial [4], the edge-Wiener index [5], and the edge-hyper-Wiener index [6]. In this paper we study the relationships between the vertex-versions and the edge-versions of the Hosoya polynomial, the Wiener index, and the edge-Wiener index of trees.

2. Preliminaries

Unless stated otherwise, the graphs considered in this paper are connected. We define $d(x, y)$ to be the distance between vertices $u, v \in V(G)$. The distance $d(e, f)$ between edges e and f of graph G is defined as the distance between vertices e and f in the line graph $L(G)$.

Email addresses: niko.tratnik1@um.si (Niko Tratnik), petra.zigert@um.si (Petra Žigert Pleteršek)

If G is a connected graph with n vertices, and if $d(G,k)$ is the number of (unordered) pairs of its vertices that are at distance k, then the *Hosoya polynomial* of G is defined as

$$H(G,x) = \sum_{k\geq 0} d(G,k)\, x^k.$$

Note that $d(G,0) = n$. Similarly, if $d_e(G,k)$ is the number of (unordered) pairs of edges that are at distance k, then the *edge-Hosoya polynomial* of G is defined as

$$H_e(G,x) = \sum_{k\geq 0} d_e(G,k)\, x^k.$$

Obviously, for any connected graph G it holds $H_e(G,x) = H(L(G),x)$.

The *Wiener index* and the *edge-Wiener index* of a connected graph G are defined in the following way:

$$W(G) = \sum_{\{u,v\}\subseteq V(G)} d(u,v), \qquad W_e(G) = \sum_{\{e,f\}\subseteq E(G)} d(e,f).$$

It is easy to see that $W_e(G) = W(L(G))$. The main property of the Hosoya polynomial and the edge-Hososya polynomial, that makes them interesting in chemistry, follows directly from the definitions (see also [7]):

$$W(G) = H'(G,1), \qquad W_e(G) = H'_e(G,1). \tag{1}$$

The *hyper-Wiener index* and the *edge-hyper-Wiener index* of a connected graph G are defined as:

$$WW(G) = \frac{1}{2}\sum_{\{u,v\}\subseteq V(G)} d(u,v) + \frac{1}{2}\sum_{\{u,v\}\subseteq V(G)} d^2(u,v),$$

$$WW_e(G) = \frac{1}{2}\sum_{\{e,f\}\subseteq E(G)} d(e,f) + \frac{1}{2}\sum_{\{e,f\}\subseteq E(G)} d^2(e,f).$$

Again, it holds $WW_e(G) = WW(L(G))$. Moreover, the following relationship was proved in [8] for any connected graph G:

$$WW(G) = H'(G,1) + \frac{1}{2}H''(G,1). \tag{2}$$

3. The edge-Hosoya polynomial of trees

In this section we first show how the edge-hyper-Wiener index of an arbitrary connected graph can be calculated from the edge-Hosoya polynomial.

Theorem 3.1. *Let G be a connected graph. Then*

$$WW_e(G) = H'_e(G,1) + \frac{1}{2}H''_e(G,1).$$

Proof. Using Equation 2 we obtain

$$WW_e(G) = WW(L(G)) = H'(L(G),1) + \frac{1}{2}H''(L(G),1) = H'_e(G,1) + \frac{1}{2}H''_e(G,1)$$

and the proof is complete. □

The following theorem is the main result of this paper.

Theorem 3.2. *Let T be a tree. Then*

$$H_e(T,x) = \frac{1}{x}H(T,x) - \frac{|V(T)|}{x}. \tag{3}$$

.

Proof. It suffices to prove that

$$H(T,x) = xH_e(T,x) + |V(T)|.$$

Let V_k be the set of all (unordered) pairs of vertices of T that are at distance k and let E_k be the set of all (unordered) pairs of edges of T that are at distance k, where $k \geq 0$. That means

$$V_k = \{\{x,y\} \mid x,y \in V(T),\ d(x,y) = k\},$$

$$E_k = \{\{e,f\} \mid e,f \in E(T),\ d(e,f) = k\}.$$

We first show that for any $k \geq 1$ there exists a bijective function $F : V_k \to E_{k-1}$. To define F, let $k \geq 1$ and let $x,y \in V(T)$ such that $d(x,y) = k$. Furthermore, let P be the unique path in T connecting x and y. Obviously, $d(x,y) = |E(P)| = k$. We define e_x to be the edge of P which has x for one end-vertex. Similarly, e_y is the edge of P which has y for one end-vertex. It is easy to see that $d(e_x,e_y) = k-1$. With this notation we can define

$$F(\{x,y\}) = \{e_x,e_y\}$$

for every $\{x,y\} \in V_k$. Obviously, F is a well-defined function.

To show that F is injective, let $\{x,y\},\{a,b\} \in V_k$, $k \geq 1$, and suppose $F(\{x,y\}) = F(\{a,b\})$. It follows that $\{e_x,e_y\} = \{e_a,e_b\}$ and without loss of generality we can assume $e_x = e_a$ and $e_y = e_b$. If $x = a$, we also get $y = b$, since otherwise $e_y \neq e_b$. Therefore, $\{x,y\} = \{a,b\}$. If $x \neq a$, it follows that $x = b$ and $y = a$. Again, $\{x,y\} = \{a,b\}$ and we are done.

To show that F is surjective, we take $\{e,f\} \in E_{k-1}$. Let x be the end-vertex of e and y the end-vertex of f such that $d(x,y) = d(e,f) + 1 = k$. Obviously, x and y are uniquely defined. It is easy to see that $F(\{x,y\}) = \{e,f\}$.

We have shown that for every $k \geq 1$ it holds $d(T,k) = |V_k| = |E_{k-1}| = d_e(T,k-1)$. It is also obvious that $d(T,0) = |V(T)|$. Hence, polynomials $H(T,x)$ and $xH_e(T,x) + |V(T)|$ have the same coefficients. Therefore, Equation 3 it true and the proof is complete. □

As a corollary we can now express the edge-Wiener index and the edge-hyper-Wiener index of trees with the Wiener index and the hyper-Wiener index.

Corollary 3.3. *Let T be a tree. Then*

$$W_e(T) = W(T) - \binom{|V(T)|}{2}$$

and

$$WW_e(T) = WW(T) - W(T).$$

Proof. First we notice that if G is a graph, then

$$H(G,1) = \sum_{k\geq 0} d(G,k) = \binom{|V(G)|}{2} + |V(G)|. \tag{4}$$

After differentiating Equation 3 we obtain

$$H'_e(T,x) = \frac{H'(T,x)x - H(T,x) + |V(T)|}{x^2} \tag{5}$$

and

$$H''_e(T,x) = \frac{H''(T,x)x^3 - 2H'(T,x)x^2 + 2H(T,x)x - 2x|V(T)|}{x^4}. \tag{6}$$

Using Equation 5 and Equation 4 it follows,

$$\begin{aligned} W_e(T) &= H'_e(T,1) \\ &= H'(T,1) - H(T,1) + |V(T)| \\ &= W(T) - \binom{|V(T)|}{2}. \end{aligned}$$

Finally, Theorem 3.1, Equation 5, Equation 6, and Equation 2 imply

$$\begin{aligned} WW_e(T) &= H'_e(T,1) + \frac{1}{2}H''_e(T,1) \\ &= H'(T,1) - H(T,1) + |V(T)| \\ &+ \frac{1}{2}H''(T,1) - H'(T,1) + H(T,1) - |V(T)| \\ &= WW(T) - W(T). \end{aligned}$$

□

4. The edge-Hosoya polynomial of dendrimers

Dendrimers are highly regular trees, which are of interest to chemists, since they represent repetitively branched molecules. In this section we compute the edge-Hosoya polynomial, the edge-Wiener index and the edge-hyper-Wiener index of regular dendrimers.

In particular, $T_{k,d}$ stands for the k-th *regular dendrimer* of degree d. For any $d \geq 3$, $T_{0,d}$ is the one-vertex graph and $T_{1,d}$ is the star with $d+1$ vertices. Then for any $k \geq 2$ and $d \geq 3$, the tree $T_{k,d}$ is obtained by attaching $d-1$ new vertices of degree one to the vertices of degree one of $T_{k-1,d}$. For an example see Figure 1. The parameter k corresponds to what in dendrimer chemistry is called "number of generations" [9].

In [10] the Wiener polynomial $W(G,x)$ of a graph G was considered and the definition of this polynomial is slightly different from the definition of the Hosoya polynomial, such that $H(G,x) = W(G,x) + |V(G)|$. Hence, from Equation 3 it follows

$$H_e(G,x) = \frac{1}{x}W(G,x).$$



Figure 1: Regular dendrimer $T_{2,3}$.

Therefore, to compute the edge-Hosoya polynomial we can use this formula and the result regarding the Wiener polynomial of a regular dendrimer in [10]. After changing some labels we obtain

$$H_e(T_{k,d},x)=\sum_{i=0}^{k-1}(d-1)^{2i}d\frac{(d-1)^{k-i}-1}{d-2}x^{2i}+\sum_{i=0}^{k-1}(d-1)^{2i}\binom{d}{2}\cdot\left(d\frac{(d-1)^{k-i-1}-1}{d-2}+1\right)x^{2i+1}.$$

It follows from Equation 1 and Theorem 3.1 that the edge-Wiener index and the edge-hyper-Wiener index can be easily computed from the derivatives of the edge-Hosoya polynomial. Therefore, we obtain

$$W_e(T_{k,d})=\frac{d\Big(2-2d+(d-1)^k(d^2+4d-4)+(d-1)^{2k}(2-d(d+2)+2(d-2)dk)\Big)}{2(d-2)^3}$$

and

$$\begin{aligned}WW_e(T_{k,d}) &= d\frac{2(d-1)+(d-1)^k\left(4-5d^2\right)}{2(d-2)^4}\\ &+ d\frac{(d-1)^{2k}\Big(-2-8k+d\left(-2+5d+16k-d(d+4)k+2(d-2)^2k^2\right)\Big)}{2(d-2)^4}.\end{aligned}$$

Since the Wiener index and the hyper-Wiener index of regular dendrimers are already known (see [9, 10, 11]), the edge-Wiener index and the edge-hyper-Wiener index could also be computed in terms of Corollary 3.3.

Acknowledgment

Supported in part by the Ministry of Science of Slovenia under grant $P1-0297$.

References

[1] H. Wiener, Structural determination of paraffin boiling points, J. Amer. Chem. Soc. 69 (1947) 17–20.

[2] H. Hosoya, On some counting polynomials in chemistry, Discrete Appl. Math. 19 (1988) 239–257.

[3] M. Randić, Novel molecular descriptor for structure-property studies, Chem. Phys. Lett. 211 (1993) 478–483.

[4] A. Behmaram, H. Yousefi-Azari, A. R. Ashrafi, Some new results on distance-based polynomials, MATCH Commun. Math. Comput. Chem. 65 (2011) 39–50.

[5] A. Iranmanesh, I. Gutman, O. Khormali, A. Mahmiani, The edge versions of Wiener index, MATCH Commun. Math. Comput. Chem. 61 (2009) 663–672.

[6] A. Iranmanesh, A. S. Kafrani, O. Khormali, A new version of hyper-Wiener index, MATCH Commun. Math. Comput. Chem. 65 (2011) 113–122.

[7] M. Petkovšek, N. Tratnik, P. Žigert Pleteršek, The edge-Hosoya polynomial of benzenoid chains, preprint.

[8] G. G. Cash, Relationship between the hosoya polynomial and the hyper-wiener index, Appl. Math. Lett. 15 (2002) 893–895.

[9] I. Gutman, Y.-N. Yeh, S.-L. Lee, J.-C. Chen, Wiener numbers of dendrimers, MATCH Commun. Math. Comput. Chem. 30 (1994) 103–115.

[10] B. E. Sagan, Y.-N. Yeh, P. Zhang, The Wiener polynomial of a graph, Int. J. Quantum Chem. 60 (1996) 959–969.

[11] M. V. Diudea, B. Parv, Molecular topology. 25. Hyper-Wiener index of dendrimers, J. Chem. Inf. Comput. Sci. 35 (1995) 1015–1018.